

<b>F.F. Wirakusumah</b> <b>R. Brand</b> <b>J.B. Gravenhorst</b>	<b>A multivariate analysis of risk factors influencing the cesarean section rate in two university hospitals in Leiden (the Netherlands) and Bandung (Indonesia)</b> During ten year period the factors that played an important role in the decision to perform a cesarean section were more or less the same in two very different countries.	<b>38</b>
<b>A.K. Janmeja</b> <b>B. Raj</b> <b>N.K. Jhamb</b> <b>B. Arora</b>	<b>Clinical trial of 4 month chemotherapy regimen in tuberculous pleural Effusion</b> Four month short course chemotherapy consisting of 2 SHZR/2HR was succesful in the treatment of tuberculous pleural effusion.	<b>45</b>
<b>E.A. Soepardi</b>	<b>Chest X-ray examination of aspirated peanut in children : is it important ?</b> Chest X-ray examination is not suggested in aspirated peanut in children before 24 hours of lodgement.	<b>49</b>

The Editorial Board wishes to apologize for publishing the following article in the *Medical Journal of Indonesia*, vol. 6 # 4, October-December 1997 edition.

"Food and social background of nasopharyngeal cancer patients in Jakarta" by A. Roezin.

This article is a duplication of an article written by the same author under the same title, published in the *Asean Otolaryngology-Head and Neck Surgery Journal* vol.1 # 1, January-March 1997 edition.