Corrigendum: Perception of the Zika virus infection and its influence on Zika prevention practices by pregnant women at the Region 5 Health Promotion Center in Thailand

Issara Siramaneerat

Department of Social Science, Faculty of Liberal Arts, Rajamangala University of Technology Thanyaburi, Thailand

Correction to: Medical Journal of Indonesia. 2018;27(3):201–8 https://doi.org/10.13-181/mji.v27i3.2424

In the previous published article,¹ there were three errors.

The first error was published on page 201, in the results section of the abstract. the result section from the abstract, the following text appears: "The findings indicated that 5 of 12 factors could significantly predict Zika prevention practices of pregnant women at the Health Promotion Center Region 5 in Thailand: education, smoking behavior, check-up status during pregnancy, perception of susceptibility, and perception of benefit." The text was incorrect and should have read: "The findings indicated that 4 of 11 factors could significantly predict Zika prevention practices of pregnant women at the Health Promotion Center Region 5 in Thailand: education, check-up status during pregnancy, perception of susceptibility, and perception of benefit."

The second error was published on page 204, left column, first paragraph. In the first paragraph, the following text appears: "The respondents were classified into six age categories: 18–20 years, 21–25 years, 26–30 years, 31–35 years, 36–40 years, and >40 years old." The text was incorrect and should have read: "The respondents were classified into five age categories: 18–20 years, 21–25 years, 26–30 years, 31–35 years and ≥36 years old."

The third error was published on page 205, in table 1, subheading age. It should have read as presented below (Table 1).

REFERENCES

1. Siramaneerat I. Perception of the Zika virus infection and its influence on Zika prevention practices by pregnant women at the Region 5 Health Promotion Center in Thailand. Med J Indones. 2018;27(3):201–8.

Characteristics Mean±SD or n (%) Coefficient 95% CI р Age 18-20 years (Ref.) 35 (8.86%) 21-25 years 0.222 (-5.207 - 1.352)62 (15.7%) -2.111 26-30 years 91 (23.04%) -0.355 0.839 (-5.508 - 1.286)31-35 years 147 (37.22%) 1.925 0.308 (-3.778 - 3.068)0.123 ≥36 years 60 (15.18%) 6.262 (-1.785 - 5.636)

Table 1. Social determinant, perception, knowledge and preventive practice of Zika

pISSN: 0853-1773 • eISSN: 2252-8083 • https://doi.org/10.13181/mji.v27i4.3415 • Med J Indones. 2018;27:310

Corresponding author: Issara Siramaneerat

issara_s@rmutt.ac.th

Copyright @ 2018 Author. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are properly cited.