

<p>S. Widjaja S. Simon A.W. Sudoyo</p>	<p><b>Subcapsular hemorrhage of the liver after laparoscopic cholecystectomy, a case report</b> After undergoing laparoscopic cholecystectomy, a 44 year old woman complained of a dull pain, that turn out to be caused by liver subcapsular hemorrhage, which was a very unusual complication.</p>	<p><b>190</b></p>
<p>H. Abdoerrachman B. Hermani</p>	<p><b>Hyoid bone release and end-to-end anastomosis on stenosis of the trachea</b> Combination of hyoid bone release and end-to-end anastomosis with insertion of T-tube has been proven efficaceous in preventing the reoccurrence of tracheal stenosis.</p>	<p><b>193</b></p>
<p>S. Tandon K.B. Gupta</p>	<p><b>Case report. Bilateral recurrent pneumothorax: a rare complication of miliary tuberculosis</b> This case report highlights the extreme rarity of bilateral recurrent pneumothorax complicating miliary tuberculosis.</p>	<p><b>199</b></p>
<p>M. Munir</p>	<p><b>Three-Dimensional (3-D) video imaging as a tool in pre-operative evaluation of head and neck tumors</b> Three-Dimensional (3-D) video imaging gives exact condition of any abnormality in head and neck. This 3-D video imaging gives more information, documentation and ability to the surgeon to handle the abnormality.</p>	<p><b>201</b></p>
<p>A. Sharma S.N. Chugh R. Kakkar S. Kharb</p>	<p><b>Letter</b> Evaluation of oxidative stress in diabetics with vascular complications.</p>	<p><b>207</b></p>

#### ERRATA FOR MED J INDONES VOL. 8 NO. 2, 1999

We apologize for some disturbing errors in some parts of the data presentations which remained during technical editing due to time constraints.

Please, notice the following corrections or clarifications:

**1. Page 78, Table 1:**

- 1.1. the data figures under STAGING for BATCH 2 of the case group were misplaced under the control group for BATCH 1 study.
- 1.2. Row "Stage II", column "n", written as 45 should read 46.

**2. Page 92, Table 2:** The data should be replaced by the following figures:

Stage I = 14 cases (4.7%), Stage II = 6 cases (15.3%), Stage III-A = 68 cases (23.7%), Stage III-B = 116 cases (38.7%) and Stage IV = 56 cases (18.6%).

**3. Page 125, Table 3:** Two cases with "Unknown" Stage were already classified and should be included in Stage IV to make up 29 cases (12.7%).

Editor in Chief  
Medical Journal of Indonesia